

ON THE PHYSICS OF TYPE I X-RAY BURSTS ON ACCRETING NEUTRON STARS AT HIGH ACCRETION RATES

RANDALL L. COOPER AND RAMESH NARAYAN

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

Draft version February 5, 2008

ABSTRACT

We investigate the effect of the hot CNO cycle breakout reaction $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ on the occurrence of type I X-ray bursts on accreting neutron stars. For $f_{\text{tp}} \lesssim 0.1$, where f_{tp} is a dimensionless factor by which we multiply the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate of Caughlan & Fowler (1988), our model predicts that bursts should occur only for accretion rates $\dot{M}$ below a critical value $\approx 0.3\dot{M}_{\text{Edd}}$. This agrees with observations. For larger values of f_{tp} , including the standard choice $f_{\text{tp}} = 1$, the model switches to a new regime in which bursts occur all the way up to $\dot{M} \approx \dot{M}_{\text{Edd}}$. Since the latter regime disagrees with observations, we suggest that the true $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate is lower than usually assumed.

Subject headings: dense matter — nuclear reactions, nucleosynthesis, abundances — stars: neutron — X-rays: binaries — X-rays: bursts

1. INTRODUCTION

Type I X-ray bursts are thermonuclear explosions that occur on accreting neutron stars in low-mass X-ray binaries. They are triggered by thermally unstable H or He burning near the stellar surface (for reviews, see Cumming 2004; Strohmayer & Bildsten 2006). For systems with accretion rates $\dot{M} \lesssim 0.1\dot{M}_{\text{Edd}}$, where $\dot{M}_{\text{Edd}}$ denotes the mass accretion rate at which the accretion luminosity is equal to the Eddington limit, the basic physics of the burst onset is well understood to be that of the thin shell thermal instability (Schwarzschild & Härm 1965; Hansen & van Horn 1975), and theoretical models have been rather successful at reproducing the gross characteristics of burst observations in this regime (e.g., Fujimoto et al. 1981; Fushiki & Lamb 1987; Cumming & Bildsten 2000; Narayan & Heyl 2003).

This *Letter* addresses a longstanding problem that afflicts nearly all burst models when the accretion rate $\dot{M} \gtrsim 0.1\dot{M}_{\text{Edd}}$. Both simple one-zone burst models (Fujimoto et al. 1981; Paczyński 1983; Bildsten 1998; Heger et al. 2005) and sophisticated time-dependent multi-zone models (Ayasli & Joss 1982; Taam et al. 1996; Fisker et al. 2003; Heger et al. 2005) predict that bursts should occur for all $\dot{M}$ up to $\approx \dot{M}_{\text{Edd}}$. Observations, however, indicate that bursts do not occur for $\dot{M} \gtrsim 0.3\dot{M}_{\text{Edd}}$ (van Paradijs et al. 1979, 1988; Cornelisse et al. 2003; Remillard et al. 2006). Furthermore, van Paradijs et al. (1988) found that, for $0.1 \lesssim \dot{M}/\dot{M}_{\text{Edd}} \lesssim 0.3$, a significant fraction of the accreted plasma burns stably between consecutive bursts, leading to large values $\gtrsim 1000$ of the parameter α , the accretion energy released between successive bursts divided by the nuclear energy released during a burst. Most theoretical models, on the other hand, predict that nearly all of the accreted matter burns unstably during bursts, giving a nearly constant $\alpha < 100$ at all $\dot{M}$.

Narayan & Heyl (2003, hereafter NH03) developed a global linear stability analysis of the accreted plasma on the surface of a neutron star. They discovered a new regime of unstable nuclear burning for $0.1 \lesssim \dot{M}/\dot{M}_{\text{Edd}} \lesssim 0.3$ that they referred to as “delayed mixed bursts.” Their model reproduced both the occurrence of considerable stable burning preceding a burst (leading to values of $\alpha \gtrsim 1000$) and the absence

of bursts for $\dot{M} \gtrsim 0.3\dot{M}_{\text{Edd}}$, in agreement with observations. However, the complexity of their model made it difficult to understand the basic physics behind delayed mixed bursts and to identify the reasons why their results differed from those of other theoretical models at high accretion rates. To remedy this, Cooper & Narayan (2006, hereafter CN06) constructed a simple two-zone model that helped elucidate the physics of delayed mixed bursts. They showed that the competition between nuclear heating due to triple- α reactions and hot CNO cycle H burning on the one hand, and radiative cooling via photon diffusion and emission on the other hand, drives an overstability that eventually triggers a thin-shell thermal instability and hence a delayed mixed burst. They asserted that H burning via the temperature-independent hot CNO cycle, augmented by the extra seed nuclei produced from stable He burning, significantly lowers the temperature sensitivity of the total nuclear energy generation rate to such an extent as to suppress the thin-shell thermal instability for $\dot{M} \gtrsim 0.3\dot{M}_{\text{Edd}}$. Therefore, H burning via the hot CNO cycle is ultimately responsible for the lower critical $\dot{M}$ above which bursts do not occur in nature.

The above argument perhaps explains why one-zone models fail, since the models generally focus only on He burning and make large approximations with respect to H burning. But why do detailed time-dependent multi-zone models with large reaction networks also perform poorly in relation to observations? These models are much more sophisticated than the models of NH03 and CN06 and consequently ought to perform better, whereas in fact the latter models agree much better with observations. CN06 hypothesized that the time-dependent multi-zone burst models may have used too large a rate for the experimentally poorly-constrained hot CNO cycle breakout reaction $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ (Wagoner 1969; Wallace & Woosley 1981; Langanke et al. 1986; Wiescher et al. 1999; Fisker et al. 2006). We test this hypothesis. We begin in §2 with a description of the model, and we present the results of the model in §3. We discuss the results in §4, and we conclude in §5.

2. THE MODEL

We use the general-relativistic global linear stability analysis of Cooper & Narayan (2005), which is an expanded and improved version of the model of NH03, to determine the sta-

bility of nuclear burning on accreting neutron stars. We assume that matter accretes spherically onto a neutron star of gravitational mass $M = 1.4M_\odot$ and areal radius $R = 10.4$ km at a rate $\dot{M}$, where $\dot{M}$ is the rest mass accreted per unit time as measured by an observer at infinity. We set the composition of the accreted matter to be that of the Sun, such that at the neutron star surface the H mass fraction $X_{\text{out}} = 0.7$, He mass fraction $Y_{\text{out}} = 0.28$, CNO mass fraction $Z_{\text{CNO,out}} = 0.016$, and heavy element fraction $Z_{\text{out}} = 0.004$, where Z refers to all metals other than CNO. In this section, we describe the modifications to the theoretical model.

For temperatures $T \gtrsim 8 \times 10^7$ K, H burns predominantly via the hot CNO cycle $^{12}\text{C}(p,\gamma)^{13}\text{N}(p,\gamma)^{14}\text{O}(\beta^+\nu)^{14}\text{N}(p,\gamma)^{15}\text{O}(\beta^+\nu)^{15}\text{N}(p,\alpha)^{12}\text{C}$, the rate of which is determined by the slow β -decays of $^{14}\text{O}(t_{1/2} = 70.6$ s) and $^{15}\text{O}(t_{1/2} = 122$ s) (Hoyle & Fowler 1965). During the hot CNO cycle, essentially all of the CNO ions are converted to ^{14}O and ^{15}O . For the range of accretion rates we consider in this investigation, the primary breakout reaction from the hot CNO cycle into the rp-process of Wallace & Woosley (1981) is $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ (Görres et al. 1995; Hahn et al. 1996; Schatz et al. 1999). If this breakout reaction occurs, the $^{19}\text{Ne}(t_{1/2} = 17.2$ s) produced in this reaction can β -decay and return to the hot CNO cycle via $^{19}\text{Ne}(\beta^+\nu)^{19}\text{F}(p,\alpha)^{16}\text{O}(p,\gamma)^{17}\text{F}(p,\gamma)^{18}\text{Ne}(\beta^+\nu)^{18}\text{F}(p,\alpha)^{15}\text{O}$, the effect of which is only to expedite the hot CNO cycle by a factor $\lesssim 1.6$. We ignore this effect in our model. If ^{19}Ne captures a proton, however, the resulting ^{20}Na ion can never return to the hot CNO cycle (Wallace & Woosley 1981). The fate of a ^{19}Ne ion thus depends on whether or not it captures a proton before it can β -decay. Therefore, the probability that a ^{19}Ne ion is removed from the hot CNO cycle is

$$R = \frac{\rho X \lambda_{p\gamma}(^{19}\text{Ne})}{\rho X \lambda_{p\gamma}(^{19}\text{Ne}) + \lambda_{\beta^+}(^{19}\text{Ne})}, \quad (1)$$

where ρ is the density, X is the H mass fraction, $\lambda_{p\gamma}(^{19}\text{Ne})$ is the $^{19}\text{Ne}(p,\gamma)^{20}\text{Na}$ reaction rate from Caughlan & Fowler (1988, hereafter CF88) with electron screening from Dewitt et al. (1973), and $\lambda_{\beta^+}(^{19}\text{Ne}) = \ln 2/t_{1/2}$ is the ^{19}Ne β -decay rate. If ^{19}Ne captures a proton, further proton captures will ensue until the nuclear flow reaches the first waiting point of the rp-process, which we assume to be ^{24}Si (Wiescher et al. 1998). The $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ reaction is the slowest reaction of this sequence, and so its rate governs the total reaction rate of this flow. Therefore, we follow Wiescher et al. (1999) and approximate the total nuclear energy generation rate of the hot CNO cycle breakout reaction sequence by

$$\epsilon_{\text{rp}} = 24E_{\text{rp}}^* \left(\frac{Y}{4} \right) \left(\frac{Z_{\text{CNO}}}{15} \right) \rho R \lambda_{\alpha\gamma}(^{15}\text{O}) \times f_{\text{rp}}, \quad (2)$$

where $E_{\text{rp}}^* = Q_{\text{rp}}/24m_p = 1.2 \times 10^{18}$ ergs g^{-1} , Y is the He mass fraction, Z_{CNO} is the mass fraction of CNO elements, and $\lambda_{\alpha\gamma}(^{15}\text{O})$ is the $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ reaction rate from CF88, which is based on the rate derived by Langanke et al. (1986). For the temperatures $T \lesssim 6 \times 10^8$ K we consider in this work, the resonant contribution of the 4.03 MeV state in ^{19}Ne dominates the $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ reaction rate (Langanke et al. 1986). The reaction rate of this contribution is directly proportional to the α width Γ_α . However, Γ_α is very difficult to measure experimentally, and so the true $\lambda_{\alpha\gamma}(^{15}\text{O})$ is highly uncertain (for recent progress on experimental constraints, see, e.g., Tan et al. 2005; Kanungo et al. 2006). To account for

this uncertainty, we multiply the CF88 rate by a dimensionless free parameter f_{rp} such that $f_{\text{rp}} = 1$ corresponds to the usual rate of CF88. We omit the electron screening contribution to the reaction rate since the enhancement factor itself depends on the relatively unconstrained resonance strength (Salpeter & van Horn 1969; Mitler 1977). Instead, we simply absorb this contribution into f_{rp} . Equations (6), (8), (9), and (10) of Cooper & Narayan (2005) thus become

$$e^{-2\Phi/c^2} \frac{\partial}{\partial \Sigma} \left(\frac{Fr^2}{R^2} e^{2\Phi/c^2} \right) = T \frac{ds}{dt} - (\epsilon_{\text{H}} + \epsilon_{\text{He}} + \epsilon_{\text{rp}} + \epsilon_{\text{N}} - \epsilon_\nu), \quad (3)$$

$$\frac{dX}{dt} = -\frac{\epsilon_{\text{H}}}{E_{\text{H}}^*} - \frac{5}{24} \frac{\epsilon_{\text{rp}}}{E_{\text{rp}}^*}, \quad (4)$$

$$\frac{dY}{dt} = \frac{\epsilon_{\text{H}}}{E_{\text{H}}^*} - \frac{\epsilon_{\text{He}}}{E_{\text{He}}^*} - \frac{4}{24} \frac{\epsilon_{\text{rp}}}{E_{\text{rp}}^*}, \quad (5)$$

$$\frac{dZ_{\text{CNO}}}{dt} = \frac{\epsilon_{\text{He}}}{E_{\text{He}}^*} - \frac{15}{24} \frac{\epsilon_{\text{rp}}}{E_{\text{rp}}^*}. \quad (6)$$

See Cooper & Narayan (2005) for the definitions of the various symbols. Note that the factor of 5 in equation (4) accounts for the five proton captures needed to reach ^{24}Si . For a substantial portion of the accretion rates we consider in this work, the nuclear flow due to stable burning extends well beyond the relatively small number of isotopes included in our limited network (e.g., Schatz et al. 1999; Fisker et al. 2005). However, the contributions from H and He burning dominate the total nuclear energy generation rate and set the thermal profile of the accreted layer, and we calculate these contributions to high accuracy. Consequently, the thermal profiles resulting from our network compare very well with those of Schatz et al. (1999) for the range of accretion rates in which our calculations overlap. Furthermore, we are interested only in the physics of the burst onset, for which only hot CNO cycle, triple- α , and breakout reactions are important. Thus, our limited network should be adequate for our purposes.

3. RESULTS

Figure 1 shows the values of the dimensionless quantity α as a function of the Eddington-scaled accretion rate $l_{\text{acc}} \equiv \dot{M}/\dot{M}_{\text{Edd}}$ for five different values of f_{rp} , where α is defined as the accretion energy released between successive bursts divided by the nuclear energy released during a burst. We follow NH03 and call a type I X-ray burst “prompt” if $\alpha \lesssim 100$ and “delayed” if $\alpha \gg 100$. The case $f_{\text{rp}} = 0$ omits the breakout reaction sequence entirely, and so this calculation is representative of the results of NH03 and CN06. The models give prompt bursts for $l_{\text{acc}} \lesssim 0.15$ and delayed bursts over the range $l_{\text{acc}} \approx 0.15$ – 0.3 . For low values of $f_{\text{rp}} \lesssim 0.1$, the effects of breakout reactions from the hot CNO cycle on the occurrence of type I X-ray bursts are minor, and the models of NH03 and CN06 continue to provide an adequate description of the nuclear physics that precedes bursts. However, the situation changes for $f_{\text{rp}} \gtrsim 0.1$. The critical l_{acc} above which delayed mixed bursts cease now significantly decreases, and the range of accretion rates in which delayed mixed bursts occur becomes notably truncated as well. More importantly, a new burst regime appears at accretion rates close to $\dot{M}_{\text{Edd}}$, where the bursts have $\alpha < 100$ and are hence prompt. This regime of bursts near the Eddington limit was hypothesized by CN06, but it was not present in either the NH03 or CN06 models due to the exclusion of hot CNO cycle breakout reactions. The usual delayed mixed burst regime and this new




FIG. 1.— Plot shows α -values of type I X-ray bursts as a function of the Eddington-scaled accretion rate $l_{\text{acc}} \equiv \dot{M}/\dot{M}_{\text{Edd}}$ for five different values of f_{tp} , the dimensionless factor by which we multiply the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate from CF88.

prompt mixed burst regime are separated by a short range of accretion rates in which bursts do not occur. These results obtained with larger values of f_{tp} agree rather well with previous theoretical models (Fujimoto et al. 1981; Ayasli & Joss 1982; Paczyński 1983; Taam 1985; Taam et al. 1996; Bildsten 1998; Fisker et al. 2003; Heger et al. 2005), most of which predicted bursts to occur up to roughly the Eddington limit. Figure 1 thus explains the origin of the differences between those models and the models of NH03 and CN06. If the breakout reaction rate is large, say $f_{\text{tp}} \gtrsim 0.1$, the results agree with most published models (which include breakout reactions in full strength), and if the rate is small, $f_{\text{tp}} \lesssim 0.1$, the results are similar to those obtained by NH03 and CN06 (who effectively set $f_{\text{tp}} = 0$). We reiterate that the latter results agree much better with observations.

4. DISCUSSION

For $l_{\text{acc}} \lesssim 0.1$, the temperatures reached during steady-state nuclear burning in the accreted layer are too low for significant leakage out of the hot CNO cycle via breakout reactions, and so these reactions have a negligible effect on the onset of type I X-ray bursts, regardless of the true $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ cross section. However, breakout reactions should affect the nuclear flow during the burst itself, and hence they could influence the burst lightcurve and possibly also the onset of subsequent bursts (Fisker et al. 2006). CN06 showed that the delayed mixed bursts of NH03 occur when Σ_{H} , the column depth at which H is depleted via stable nuclear burning as measured from the stellar surface, is less than but close to Σ_{He} , the column depth at which He is depleted via stable nuclear burning. The hot CNO cycle breakout reaction sequence (i) eliminates seed nuclei from the hot CNO cycle which slows H burning, thereby increasing Σ_{H} , and (ii) provides an additional pathway by which He may burn, thereby decreasing Σ_{He} . Consequently, one expects the regime of delayed mixed bursts to occur at lower l_{acc} if breakout reactions are included. Furthermore, CN06 showed that the interplay between H burning via the hot CNO cycle and He burning via triple- α reactions is integral to generating the oscillations that precede delayed mixed bursts. Breakout reactions diminish this interplay by eliminating the hot CNO cycle seed nuclei and should there-

fore reduce the range of accretion rates over which delayed mixed bursts occur. Figure 1 illustrates both of these effects (compare $f_{\text{tp}} = 1$ with $f_{\text{tp}} = 0$).

For accretion rates above those at which delayed mixed bursts occur, NH03 found that nuclear burning is always stable and therefore bursts do not occur, in accord with observations (van Paradijs et al. 1979, 1988; Cornelisse et al. 2003; Remillard et al. 2006). According to their model, for $0.3 \lesssim l_{\text{acc}} \lesssim 1$, steady-state H burning via the hot CNO cycle increases the effective radiative cooling rate and thereby suppresses a He-triggered thin-shell thermal instability (CN06). However, their model did not include breakout reactions. If the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ breakout reaction rate is significant in this regime, this reaction will suppress hot CNO cycle H burning, and thus He burning via triple- α reactions will govern the total reaction rate, since the triple- α reaction rate will be the slowest rate in the nuclear flow at these high l_{acc} . The notion that the triple- α reaction rate is the slowest rate in the nuclear flow is precisely what one-zone type I X-ray burst ignition models assume (Fujimoto et al. 1981; Paczyński 1983; Bildsten 1998; Cumming & Bildsten 2000; Heger et al. 2005), and we suggest that this is the reason why the results of one-zone and time-dependent multi-zone models agree so well. All of these models predict that nuclear burning is thermally stable for temperatures $T \gtrsim 5 \times 10^8$ K, which are reached only at accretion rates $l_{\text{acc}} \gtrsim 1$ (e.g., Schatz et al. 1999). Figure 1 illustrates that, for relatively large values of f_{tp} , there is a regime of prompt bursts at accretion rates greater than the accretion rates at which delayed mixed bursts occur, and the critical accretion rate above which these prompt mixed bursts cease is roughly $\dot{M}_{\text{Edd}}$, in very good agreement with these other models. Figure 1 illustrates also that nuclear burning is stable for the small range of l_{acc} between these two regimes. These accretion rates are high enough to suppress delayed mixed bursts, but they are not high enough to cause a sufficient leakage out of hot CNO cycle H burning and trigger a thermal instability.

The model we present in this work suggests that, if the true $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ cross section is greater than approximately 0.1 of the CF88 rate, type I X-ray bursts should occur in systems with accretion rates near the Eddington limit. Observations indicate that this is not the case, since low-mass X-ray binaries with $l_{\text{acc}} \gtrsim 0.3$ generally do not exhibit bursts. This suggests that the true $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate is less than that proposed by CF88. This conclusion is complementary to that of Fisker et al. (2006), who were the first to propose that the occurrence of type I X-ray bursts is sensitive to the strength of the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate, and who found that the existence of bursts in systems with $l_{\text{acc}} \approx 0.1$ suggests a lower bound on this rate. This lower bound corresponds to $f_{\text{tp}} \approx 0.05$ in our notation. We also note that a relatively low $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate is consistent with the existence of carbon-triggered superbursts, since a low rate would increase the carbon yield resulting from stable nuclear burning (Schatz et al. 1999; Cooper et al. 2006; Fisker et al. 2006). The systems GX 17+2 and Cyg X-2 are exceptions to this empirical rule, however, for they exhibit bursts at accretion rates near the Eddington limit (e.g., Kahn & Grindlay 1984; Tawara et al. 1984; Kuulkers et al. 2002). However, it is possible that they show bursts for other reasons, e.g., they harbor mass donor stars with H-deficient envelopes (Cooper et al. 2006).

Although the resonant contribution of the 4.03 MeV state dominates the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate for the temperatures

$T \lesssim 5 \times 10^8$ K at which helium burning can possibly trigger a type I X-ray burst, other resonances such as the 4.38 MeV state contribute as well. However, these resonances contribute much less than a tenth of the 4.03 MeV contribution (e.g., Davids et al. 2003), whereas a reduction of the total $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate by a factor of only ≈ 10 is needed to stabilize nuclear burning. Therefore, it is the 4.03 MeV α width alone that determines the stability of nuclear burning at high accretion rates.

5. CONCLUSIONS

Using the global linear stability analysis of Cooper & Narayan (2005), which is an expanded and improved version of the model of NH03, we have investigated the effects of the hot CNO cycle breakout reaction $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ on the occurrence of type I X-ray bursts on accreting neutron stars at high accretion rates. For low values of $f_{\text{tp}} \lesssim 0.1$, where f_{tp} is a dimensionless factor by which we multiply the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate of CF88, the hot CNO cycle breakout reaction slightly lowers the critical accretion rate above which delayed mixed bursts occur, but otherwise the breakout reaction has little effect on the burst onset. The predictions of these models are in good agreement with observations. For $f_{\text{tp}} \gtrsim 0.1$, a new regime of prompt mixed bursts appears at accretion rates above the rates at

which delayed mixed bursts occur, and the bursts survive up to roughly the Eddington limit. The existence of this prompt mixed burst regime up to the Eddington limit is consistent with nearly all previous theoretical models. Our results support the hypothesis of CN06 that the discrepancies between the results of the global linear stability analysis of NH03 and the results of time-dependent multi-zone models with large reaction networks may be caused by the latter models assuming too large a strength for the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate at temperatures $T \lesssim 6 \times 10^8$ K. The fact that observations agree much better with the results of NH03 and CN06 implies that the true reaction rate is lower than the rate assumed in these multi-zone models. Specifically, we suggest that the true α width Γ_α of the 4.03 MeV state in ^{19}Ne is lower than the α width proposed by Langanke et al. (1986). Calculations using multi-zone models such as those of Woosley et al. (2004) and Fisker et al. (2006), but with the $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ reaction rate lowered in strength, need to be carried out to either confirm or refute this suggestion.

We thank the referee for several insightful comments and suggestions. This work was supported by NASA grant NNG04GL38G.

REFERENCES

- Ayasli, S. & Joss, P. C. 1982, *ApJ*, 256, 637
Bildsten, L. 1998, in *NATO ASIC Proc. 515: The Many Faces of Neutron Stars*, 419
Caughlan, G. R. & Fowler, W. A. 1988, *Atomic Data and Nuclear Data Tables*, 40, 283 (CF88)
Cooper, R. L., Mukhopadhyay, B., Steeghs, D., & Narayan, R. 2006, *ApJ*, 642, 443
Cooper, R. L. & Narayan, R. 2005, *ApJ*, 629, 422
—. 2006, accepted by *ApJ* (astro-ph/0605001) (CN06)
Cornelisse, R., in 't Zand, J. J. M., Verbunt, F., Kuulkers, E., Heise, J., den Hartog, P. R., Cocchi, M., Natalucci, L., Bazzano, A., & Ubertini, P. 2003, *A&A*, 405, 1033
Cumming, A. 2004, *Nuclear Physics B Proceedings Supplements*, 132, 435
Cumming, A. & Bildsten, L. 2000, *ApJ*, 544, 453
Davids, B., et al. 2003, *Phys. Rev. C*, 67, 065808
Dewitt, H. E., Graboske, H. C., & Cooper, M. S. 1973, *ApJ*, 181, 439
Fisker, J. L., Brown, E. F., Liebendörfer, M., Thielemann, F.-K., & Wiescher, M. 2005, *Nuclear Physics A*, 752, 604
Fisker, J. L., Gorres, J., Wiescher, M., & Davids, B. 2006, accepted by *ApJ* (astro-ph/0410561)
Fisker, J. L., Hix, W. R., Liebendörfer, M., & Thielemann, F.-K. 2003, *Nuclear Physics A*, 718, 614
Fujimoto, M. Y., Hanawa, T., & Miyaji, S. 1981, *ApJ*, 247, 267
Fushiki, I. & Lamb, D. Q. 1987, *ApJ*, 323, L55
Görres, J., Wiescher, M., & Thielemann, F.-K. 1995, *Phys. Rev. C*, 51, 392
Hahn, K. I., et al. 1996, *Phys. Rev. C*, 54, 1999
Hansen, C. J. & van Horn, H. M. 1975, *ApJ*, 195, 735
Heger, A., Cumming, A., & Woosley, S. E. 2005, *ApJ*, submitted (astro-ph/0511292)
Hoyle, F. & Fowler, W. A. 1965, in *Quasi-Stellar Sources and Gravitational Collapse*, 17
Kahn, S. M. & Grindlay, J. E. 1984, *ApJ*, 281, 826
Kanungo, R., et al. 2006, *Phys. Rev. C*, submitted (nucl-ex/0605033)
Kuulkers, E., Homan, J., van der Klis, M., Lewin, W. H. G., & Méndez, M. 2002, *A&A*, 382, 947
Langanke, K., Wiescher, M., Fowler, W. A., & Gorres, J. 1986, *ApJ*, 301, 629
Mittler, H. E. 1977, *ApJ*, 212, 513
Narayan, R. & Heyl, J. S. 2003, *ApJ*, 599, 419 (NH03)
Paczynski, B. 1983, *ApJ*, 264, 282
Remillard, R. A., Lin, D., Cooper, R. L., & Narayan, R. 2006, *ApJ*, 646, 407
Salpeter, E. E. & van Horn, H. M. 1969, *ApJ*, 155, 183
Schatz, H., Bildsten, L., Cumming, A., & Wiescher, M. 1999, *ApJ*, 524, 1014
Schwarzschild, M. & Härm, R. 1965, *ApJ*, 142, 855
Strohmayer, T. & Bildsten, L. 2006, in *Compact Stellar X-Ray Sources*, ed. W. H. G. Lewin and M. van der Klis (Cambridge: Cambridge Univ. Press)
Taam, R. E. 1985, *Annual Review of Nuclear and Particle Science*, 35, 1
Taam, R. E., Woosley, S. E., & Lamb, D. Q. 1996, *ApJ*, 459, 271
Tan, W. P., Görres, J., Daly, J., Couder, M., Couture, A., Lee, H. Y., Stech, E., Strandberg, E., Ugalde, C., & Wiescher, M. 2005, *Phys. Rev. C*, 72, 041302
Tawara, Y., Hirano, T., Kii, T., Matsuoka, M., & Murakami, T. 1984, *PASJ*, 36, 861
van Paradijs, J., Cominsky, L., Lewin, W. H. G., & Joss, P. C. 1979, *Nature*, 280, 375
van Paradijs, J., Penninx, W., & Lewin, W. H. G. 1988, *MNRAS*, 233, 437
Wagoner, R. V. 1969, *ApJS*, 18, 247
Wallace, R. K. & Woosley, S. E. 1981, *ApJS*, 45, 389
Wiescher, M., Görres, J., & Schatz, H. 1999, *Journal of Physics G Nuclear Physics*, 25, 133
Wiescher, M., Schatz, H., & Champagne, A. E. 1998, *Royal Society of London Philosophical Transactions Series A*, 356, 1
Woosley, S. E., Heger, A., Cumming, A., Hoffman, R. D., Pruet, J., Rauscher, T., Fisker, J. L., Schatz, H., Brown, B. A., & Wiescher, M. 2004, *ApJS*, 151, 75